



Figure 1

Figure 1 shows the temperature of 1.00 kilograms of ice ( $\text{H}_2\text{O}$ ) starting at  $-20^\circ\text{C}$  that is heated at a constant rate of 100 Joules per second (100 J/s). After about 8.6 hours, the ice has become water vapor (still  $\text{H}_2\text{O}$ !) at  $120^\circ\text{C}$ . There are five distinct regions in the curve, each of which has an arrow nearby.

1. Above each arrow *write what is happening* to the  $\text{H}_2\text{O}$  in that region. (See the example for the first region.)
2. Below each line, *write the algebra equation that is used to calculate the heat* ( $q$ ) for that region. Label the heat ( $q_2, q_3, q_4, q_5$ ) in order from left to right. (See the example for the first region.)

3. For each of the five regions, use your equation and the information provided in the figure 1 caption and in table 1 to calculate the heat for each region using the *numerical information*. Complete your calculations on the next page – be sure to show your work and include all units with each calculation. Record your results in the appropriate answer blank. Use a separate piece of paper if you need more space. Keep the paper with this worksheet.

Table 1: Key thermochemical data for water			
$C_{\text{ice}} =$	2.11	J/(g C)	$\Delta H_f =$ 333.6 J/g
$C_{\text{water}} =$	4.18	J/(g C)	$\Delta H_v =$ 2256 J/g
$C_{\text{steam}} =$	2.08	J/(g C)	$q_{\text{total}} = q_1 + q_2 + q_3 + q_4 + q_5$

Show calculations here:

- a.  $q_1 =$  \_\_\_\_\_
- b.  $q_2 =$  \_\_\_\_\_
- c.  $q_3 =$  \_\_\_\_\_
- d.  $q_4 =$  \_\_\_\_\_
- e.  $q_5 =$  \_\_\_\_\_
- f.  $q_{\text{total}} =$  \_\_\_\_\_

4. Which segment represents the largest amount of energy? Explain why.
5. Which phase (solid, liquid, or gas) takes the most energy to warm up by  $20^\circ\text{C}$ ?
- a. Explain how you can tell by looking at the graph.
- b. Explain how you can tell by looking at the values of specific heat.
6. In an experiment, 2.0 grams of NaOH are dissolved in 100 g of water. The temperature of the water goes from  $21^\circ\text{C}$  to  $25^\circ\text{C}$ .
- a. Is the reaction endothermic or exothermic?
- b. Calculate the following;
- i. mass of the solution =  $m =$
- ii. change in temperature =  $\Delta T =$
- iii. heat =  $q =$
- Given that the specific heat of the solution =  $4.18 \text{ J/g}^\circ\text{C}$ .
- iv. moles of NaOH =
- v. enthalpy for the reaction in units of  $\text{kJ/mol} = \Delta H =$